



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Smallpox in the United States, etc.—Continued.

Place,	Date.	Cases.	Deaths.	Remarks.
Washington—Continued.				
Whitman County	May 1-Aug. 31	7	2	
Yakima County	May 1-Oct. 31	13	
Total for State		110	8	
Total for State, same period, 1903.		319	11	
Wisconsin:				
Milwaukee	June 12-Dec. 17	209	
Total for State		209	
Total for State, same period, 1903.		1,732	12	
Grand total		6,598	197	
Grand total, same period, 1903.		13,739	606	

Plague in the United States, as reported to the Surgeon-General Public Health and Marine-Hospital Service, June 25 to December 30, 1904—No cases—Last case reported March 1, 1904.

California. Summary: Calendar year 1900, 22 cases, 22 deaths; 1901, 30 cases, 25 deaths; 1902, 41 cases, 41 deaths; 1903, 17 cases, 17 deaths; six months ended July 1, 1904, 8 deaths and 1 recovery.

Yellow fever in the United States, as reported to the Surgeon-General Public Health and Marine-Hospital Service, June 25 to December 30, 1904—No cases—Last case reported March 18, 1904.

Six cases reported at Laredo, Tex. (including 1 imported from Minera), during the six months ended July 1, 1904.

Weekly mortality table, cities of the United States.

Cities.	Week ended—	Population, United States census of 1900.	Total deaths from all causes.	Deaths from—									
				Tuberculosis.	Yellow fever.	Smallpox.	Varioloid.	Cholera.	Typhus fever.	Enteric fever.	Scarlet fever.	Diphtheria.	Measles.
Altoona, Pa.	Dec. 17	38,973	1									
Berkeley, Cal.	do	13,214	4	1									
Boston, Mass.	do	560,892	209	22					3	1	5		1
Brockton, Mass.	do	40,063	13	1									
Cambridge, Mass.	do	91,886	20	1									
Camden, N. J.	do	75,935	1										
Charleston, S. C.	Dec. 10	55,807	30	2									
Do	Dec. 17	55,807	17	3									
Chelsea, Mass.	do	34,072	20	3									
Cincinnati, Ohio.	Dec. 16	325,902	137	18					2	2		2	
Danville, Ill.	Dec. 17	16,354	9	3					5				
Dayton, Ohio.	do	85,333	26	2								1	
Dunkirk, N. Y.	do	11,616	6	1									
Elmira, N. Y.	do	35,672	11	1									
Erie, Pa.	Dec. 16	52,733	21	2									
Everett, Mass.	Dec. 17	24,336	2										
Galesburg, Ill.	do	18,607	7										
Hamilton, Ohio.	Nov. 5	23,914	4	1									
Do	Nov. 12	23,914	12	2									1
Do	Nov. 19	23,914	13	1									
Do	Nov. 26	23,914	5										
Hydepark, Mass.	Dec. 17	13,244	3										
Jacksonville, Fla.	do	28,429	11	2									
Lawrence, Mass.	do	62,559	25	1					1				
Lexington, Ky.	do	26,369	11	1					1				
Los Angeles, Cal.	Dec. 10	102,479	69	12									
Louisville, Ky.	Dec. 15	204,731	84	6								2	

